



Shoulder Instability – Arthroscopic / Open Stabilisation

Introduction

The shoulder joint is designed to give a large amount of movement. Therefore support from the ligaments and muscles is essential.



When the shoulder dislocates the ligaments can be torn or stretched and in some cases need to be repaired.



The Operation

The operation is done by 'key hole surgery'; usually through two or three 5mm puncture wounds. The operation involves repairing the over-stretched or torn ligaments deep around the shoulder joint. The repair involves stitching the torn or stretched ligaments back onto its attachment to the socket of the shoulder blade (Glenoid). This is done using tiny anchors with sutures attached to them. The repair should be protected until healing take place (for initial healing - 6 weeks).

If not suitable, the surgeon may perform open stabilisation. (Rare - less than 2%).



General Advice

You will usually be in hospital either for a day or overnight. The operation is performed under general anaesthetic and a nerve block in your neck or upper chest that will make your arm 'dead' for 8-12 hours after surgery. This is for post operative pain relief. After this the shoulder may well be sore and you will be given painkillers to help this. Ice packs may also help reduce pain. Wrap crushed ice or frozen peas in a damp, cold cloth and place on the shoulder for up to 15 minutes.

You will return from theatre wearing a sling. Your arm will remain in a special sling for 6 weeks. This means that you will be unable to use the arm throughout this time and you will be unable to return to work. You will not be allowed to drive for 6-8 weeks after surgery.

The shoulder must remain immobilised with a sling and a body belt (underneath the clothes) for 3 weeks. At your follow-up appointment the body belt will be removed and you will be allowed to wear the sling on top of your clothes.

You will be expected to remove the sling for exercises only. Your physiotherapist will advise you of these.

A doctor/physiotherapist will see you prior to discharge and you will be taught exercises to do and given further advice to guide you through your recovery.

The length of time that you will be off work will depend on your job but expect a minimum of 6 weeks.

The wounds:

Arthroscopic (keyhole) stabilisation:

This keyhole operation is usually done through two or three 4mm puncture wounds. There may be no stitches (may be only one absorbable stitch in the front wound) only small sticking plaster strips over the wounds. These should be kept dry until healed. This usually takes 5 to 7 days.

Open stabilisation:

There is an incision at the front of the armpit within the natural skin crease. The stitch is dissolvable but is usually removed at 3 weeks. Keep the wound dry until it is well healed.

Sport / Leisure Activities

Your physiotherapist and surgeon will advise you when it is safe to resume your leisure activities. However, you **will NOT be allowed** to return to contact sports before 6 months from surgery.

Complications

As with all surgery there is a risk of some complications. These are rare, but you should be aware of them before your operation.

They include:

1. Complications relating to the anaesthetic.
2. Infection.
 - Failure to achieve a successful result.
 - A need to redo the surgery.
 - Injury to the nerves or blood vessels around the shoulder.
 - Fracture
 - Prolonged stiffness and or pain.
 - Implant failure
 - A further dislocation of the shoulder (less than 5%).

If you require further information please discuss with the doctors either in clinic or on admission.

Exercises

The shoulder must remain immobilised with a sling and a body belt (underneath the clothes) for 3 weeks. At your follow-up appointment the body belt will be removed and you will be allowed to wear the sling on top of your clothes.

You will perform first only elbow, wrist and hand exercises.

You may well be expected to perform the following exercises when you leave hospital. Please check with your physiotherapist before commencing.

Keep your arm in the sling and move your hand up and down at the wrist.

With your arm out of the sling bend and straighten the elbow

With your arm in the sling and the elbow bent at your side, turn the hand to face the ceiling and then the ground.

With arm in the sling regularly shrug shoulders up and down and circle forwards and backwards

Repeat these exercises four times per day. The number you should perform at each session will be recorded for you by your physiotherapist.

Continue these exercises until otherwise advised by your physiotherapist.

If you require further information please discuss this with the doctors either in clinic or on admission.



Appointments:

Private:

Professor Ofer Levy, MD MCh(Orth) FRCS

Practice Manager: Cherie Emmitt
Telephone: 07367 582777

e-mail: bm@readingshoulderunit.com

NHS:

**Reading Shoulder Unit
NHS Choose & Book -**

The Berkshire Independent Hospital
Swallows Croft, Wensley Road,
Reading, RG1 6UZ

Secretary: Helen Henly
Telephone: 0118 902 8109

e-mail: Helen.Henly@ramsayhealth.co.uk

**Reading Shoulder Unit
NHS Choose & Book -**

Spire Dunedin Hospital
16 Bath Road,
Reading, RG1 6UZ

Secretary: Karen Dawson
Telephone: 0118 955 3496

email: karen.dawson2@spirehealthcare.com

For patient testimonials see website:

<http://www.readingshoulderunit.com/patient-information/feedback>

Industry Relationships

Reading Shoulder Unit surgeons and scientists may collaborate with the pharmaceutical or medical device industries to help develop medical breakthroughs or provide medical expertise or education. Reading Shoulder Unit strives to make scientific advances that will benefit patient care and support outside relationships that promise public benefit. In order for the discoveries of Reading Shoulder Unit surgeons' and scientists' laboratories and investigations to benefit the public, these discoveries must be commercialised in partnership with industry. As experts in their fields, Reading Shoulder Unit surgeons and scientists are often sought after by industry to consult, provide expertise and education. To assure professional and commercial integrity in such matters, Reading Shoulder Unit maintains a program that reviews these collaborations and, when appropriate, puts measures in place to minimise bias that may result from ties to industry. In publicly disclosing this information, the Reading Shoulder Unit tries to provide information as accurately as possible about its surgeons' and scientists' connections with industry.

Professor Levy has a professional relationship with the company that makes the Verso prosthesis. He helped design it and acts as their Chief Medical Advisor and Company Secretary. However, he does not receive any money, royalties, or financial rewards when this device is used. He holds shares in the manufacturing company in lieu of granting the patent. He is telling you this so you have full transparency. He and his team believe that the treatment offered is the best for your specific medical needs, but you are completely free to ask for an alternative or a second opinion

